

BRAZILIAN BUTT LIFT POST-OPERATIVE INSTRUCTIONS

Supplies to have at home before surgery:

- Prescription medications
- Acetaminophen (Tylenol) and ibuprofen (Advil)
- Dimenhydrinate (Gravol) (anti-nausea medication)
- Constipation medications (examples: Senokot, Restoralax)
- Garbage bag / old towels (to protect the car seats on your ride home from surgery)
- Sanitary pads

Activity:

- After surgery you will have absorbent sanitary pads present on some of your liposuction incisions, and you will be wearing a compression garment
- The pads can be changed as needed over the next 48 hours. It is normal for the liposuction incisions to drain during this time.
- The garment is to be worn for 23 hours per day for four weeks after surgery. Each day you can take two 30 min breaks where you are not wearing the binder.
- Each hour, get up and walk around the house
- When resting, laying on your stomach is optimal
- Avoid sitting directly on your buttocks, although you can sit to eat, use the bathroom and travel short distances
- Take 10 deep breaths every hour while awake
- Showering can be done 48 hours after surgery
- Avoid excessive physical activity or lifting objects greater than 10 lbs for 4 weeks after surgery
- Don't sleep with pets in bed for the first two weeks after surgery
- Avoid smoking or exposure to second-hand smoke
- No baths, swimming, or hot tubs for four weeks after surgery

Medications:

- Ibuprofen (Advil) 400 mg is to be taken 3 times per day: breakfast, lunch and dinner with food. **NEXT DOSE:**
- Acetaminophen (Tylenol) 500 mg is to be taken 4 times per day: breakfast, lunch, dinner and bedtime. **NEXT DOSE:**

- Narcotic prescription hydromorphone (Dilaudid) or Tylenol #3 is to be taken every 4 to 6 hours, ONLY as needed. **NEXT DOSE:**
- Antibiotic (cephalexin or clindamycin) is to be started the morning after surgery until completed. If you were not prescribed antibiotics, you do not require them for your surgery.
- Constipation medications can be taken according to package instructions if no bowel movement for 2 days after surgery
- Dimenhydrinate (Gravol) may be taken according to package instructions for post-operative nausea and vomiting

First Post-operative Appointment:

- Your sutures will be removed at this appointment
- Swelling, bruising and some degree of asymmetry is expected at this appointment

Wound Care:

- Any dried blood along the incision lines can be safely removed with the use of a Q-tip dipped in hydrogen peroxide or warm water. This can be done twice a day, followed with a thin application of Polysporin to the incision line.
- Do not continue to apply Polysporin to the incision line daily past 1 week. This can result in a rash or allergic reaction.

Recovery:

- Massage of the liposuction donor areas can be done at 2 weeks after surgery, but do not massage your buttocks at this time.
- Sunscreen is extremely important. The use of SPF 45 or higher along the incision lines will prevent pigment changes due to sun exposure. This can be started 2-3 weeks after surgery. This needs to be done while wearing a bathing suit or light clothing.

RED FLAG SYMPTOMS:

If any of the following symptoms develop post-operatively, please contact the clinic immediately.

- Sudden increase in swelling at the surgical site
- Increasing pain, redness, foul odour or drainage from the incision
- Sudden increase in swelling in the cheek, jawline or neck
- Fever exceeding 38.3°C

- Persistent bleeding from an incision
- Sudden onset of chest pain or shortness of breath (For chest pain lasting longer than 5 minutes, call 911)
- Pain or swelling of the leg
- Persistent dizziness or loss of consciousness
- Coughing up blood or bloody stool

If you develop any of these symptoms and are unable to reach the clinic or Dr. MacArthur, present to the nearest Emergency Room.

Do not hesitate to call with any questions or concerns.

Clinic business hours (9AM to 4 PM): 613-595-1880

After hours: 613-219-9636 (Dr. MacArthur Cell)

Patient signature: _____

Date: _____

RN signature: _____

Date: _____